Glossary

Blinding (masking): in an experimental study, refers to whether patients, clinicians providing an intervention, people assessing outcomes, and/or data analysts were aware or unaware of the group to which patients were assigned. In the design section of Evidence-Based Nursing abstracts of treatment studies, the study is identified as *blinded*, with specification of who was blinded; unblinded, if all parties were aware of patients' group assignments; or blinded (unclear) if the authors did not report or provide us with an indication of who was aware or unaware of patients' group assignments.

Concealment of randomisation: concealment of randomisation is specified in the design section of Evidence-Based Nursing abstracts of treatment studies as follows: allocation concealed (deemed to have taken adequate measures to conceal allocation to study group assignments from those responsible for assessing patients for entry in the trial [ie, central randomisation; sequentially numbered, opaque, sealed envelopes; sealed envelopes from a closed bag; numbered or coded bottles or containers; drugs prepared by the pharmacy; or other descriptions that contain elements convincing of concealment]); allocation not concealed (deemed to have not taken adequate measures to conceal allocation to study group assignments from those responsible for assessing patients for entry in the trial [ie, no concealment procedure was undertaken, sealed envelopes that were not opaque or were not sequentially numbered, or other descriptions that contained elements not convincing of concealment]); unclear allocation concealment (the authors did not report or provide a description of an allocation concealment approach that allowed for the classification as concealed or not concealed).

Confidence interval (CI): quantifies the uncertainty in measurement; usually reported as 95% CI, which is the range of values within which we can be 95% sure that the true value for the whole population lies.

Constant comparison¹: a procedure used in qualitative research wherein newly collected data are compared in an ongoing fashion with data obtained earlier, to refine theoretically relevant categories.

Cluster randomisation²: randomisation of groups of people rather than individuals; this approach is often used to avoid "contamination" when the way in which people in one group are treated or assessed is likely to modify the treatment or assessment of people in other groups.

Crossover trial: a method of comparing 2 interventions in which patients are switched to the alternative intervention after a specified period of time.

Data saturation (saturation, redundancy)1: process of collecting data in a qualitative research study to the point where no new themes are generated.

Fixed effects model³: gives a summary estimate of the magnitude of effect in meta-analysis. It takes into account within-study variation but not between-study variation and hence is usually not used if there is significant heterogeneity.

Heterogeneity³: the degree to which the effect estimates of individual studies in a meta-analysis differ significantly.

Intention to treat analysis (ITT): all patients are analysed in the groups to which they were randomised, even if they failed to complete the intervention or received the wrong intervention.

Logistic regression⁴: a statistical technique that determines the probability of a dependent variable (outcome) occurring when the independent (explanatory) variables are present or absent when the outcome is a dichotomous (binary) variable. It determines whether a model that includes the variable(s) explains more about the outcome variable than a model that does not include the variable(s).

Number needed to treat (NNT): number of patients who need to be treated to prevent 1 additional negative event (or to promote 1 additional positive event); this is calculated as 1/absolute risk reduction (rounded to the next whole number), accompanied by the 95% confidence interval.

Number needed to harm (NNH)5: number of patients who, if they received the experimental treatment, would lead to 1 additional person being harmed compared with patients who receive the control treatment; this is calculated as 1/absolute risk increase (rounded to the next whole number), accompanied by the 95% confidence interval.

Odds ratio (OR): describes the odds of a patient in the experimental group having an event divided by the odds of a patient in the control group having the event or the odds that a patient was exposed to a given risk factor divided by the odds that a control patient was exposed to the risk factor.

Random effects model³: gives a summary estimate of the magnitude of effect in meta-analysis. It takes into account both within-study and between-study variance and gives a wider confidence interval to the estimate than a fixed effects model if there is significant between-study variation.

Relative benefit increase (RBI): the proportional increase in the rates of good events between experimental and control participants; it is reported as a percentage (%).

Relative risk increase (RRI): the proportional increase in bad outcomes between experimental and control participants; it is reported as a percentage (%).

Relative risk reduction (RRR): the proportional reduction in bad outcomes between experimental and control participants; it is reported as a percentage (%).

Receiver operating characteristic (ROC) curve⁶: an analysis used to assess the clinical usefulness of a diagnostic or screening test. It yields a score that has the highest rates of both sensitivity and specificity with respect to a diagnosis-that is, a score that will give the maximum rate of accurate classifications.

Sensitivity⁵: a measure of a diagnostic test's ability to correctly detect a disorder when it is present in a sample of people.

Specificity⁵: a measure of a diagnostic test's ability to correctly identify the absence of a disorder in a sample of people who do not have the disorder.

Weighted mean difference³: in a meta-analysis, used to combine outcomes measured on continuous scales (eg, height), assuming that all trials measured the outcome on the same scale; the mean, standard deviation and sample size of each group are known, and weight given to each trial is determined by the precision of its estimate of effect.

- Polit DF, Hungler BP. Essentials of nursing research: methods, appraisal, and utilization. Fourth edition. Philadelphia: Lippincott, 1997. 1
- 9 Jadad AR. Randomised controlled trials. London: BMJ Books, 1998.
- Clarke M, Oxman AD, editors. Glossary. Cochrane reviewers' handbook 4.1.2 (updated March 2001). In: *Cochrane Library*, Oxford: Update Software. Updated quarterly.
- Dawson-Saunders B, Trapp RG. Basic and clinical biostatistics. Norwalk: Appleton and Lange, 1994. 4
- Sackett DL, Haynes RB, Guyatt GH, et al. Clinical epidemiology: basic science for clinical medicine. Second edition. Boston: Little, Brown and Company, 1991.
- RA, Cavalieri TA, Leonard DM, et al. Gen Hosp Psychiatry 6 Steer 1999:21:106-11